

## Sustainable, efficient livestock production with high biodiversity and good welfare for animals

D. M. Broom, F. A. Galindo and E. Murgueitio

*Proc. R. Soc. B* 2013 **280**, 20132025, published 25 September 2013

---

### References

This article cites 44 articles, 4 of which can be accessed free  
<http://rsbp.royalsocietypublishing.org/content/280/1771/20132025.full.html#ref-list-1>

### Subject collections

Articles on similar topics can be found in the following collections

[environmental science](#) (230 articles)

### Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right-hand corner of the article or click [here](#)

**Review**



**Cite this article:** Broom DM, Galindo FA, Murgueitio E. 2013 Sustainable, efficient livestock production with high biodiversity and good welfare for animals. *Proc R Soc B* 280: 20132025.

<http://dx.doi.org/10.1098/rspb.2013.2025>

Received: 2 August 2013

Accepted: 4 September 2013

**Subject Areas:**

environmental science

**Keywords:**

sustainability, silvopastoral, livestock, biodiversity, animal welfare, conservation

**Author for correspondence:**

D. M. Broom

e-mail: [dmb16@cam.ac.uk](mailto:dmb16@cam.ac.uk)

# Sustainable, efficient livestock production with high biodiversity and good welfare for animals

D. M. Broom<sup>1</sup>, F. A. Galindo<sup>1,2</sup> and E. Murgueitio<sup>3</sup>

<sup>1</sup>Centre for Animal Welfare and Anthrozoology, Department of Veterinary Medicine, University of Cambridge, Madingley Road, Cambridge CB3 0ES, UK

<sup>2</sup>Departamento de Etología, Fauna Silvestre y Animales de Laboratorio (DEFSAL), Facultad de Medicina Veterinaria y Zootecnia (FMVZ), Universidad Nacional Autónoma de México (UNAM), Ciudad Universitaria, 04510 Mexico City, Mexico

<sup>3</sup>Fundación CIPAV, Cra. 25 No. 6-62, Cali, Colombia

What is the future for livestock agriculture in the world? Consumers have concerns about sustainability but many widely used livestock production methods do not satisfy consumers' requirements for a sustainable system. However, production can be sustainable, occurring in environments that supply the needs of the animals resulting in good welfare, allow coexistence with a wide diversity of organisms native to the area, minimize carbon footprint and provide a fair lifestyle for the people working there. Conservation need not just involve tiny islands of natural vegetation in a barren world of agriculture, as there can be great increases in biodiversity in farmed areas. Herbivores, especially ruminants that consume materials inedible by humans, are important for human food in the future. However, their diet should not be just ground-level plants. Silvopastoral systems, pastures with shrubs and trees as well as herbage, are described which are normally more productive than pasture alone. When compared with widely used livestock production systems, silvopastoral systems can provide efficient feed conversion, higher biodiversity, enhanced connectivity between habitat patches and better animal welfare, so they can replace existing systems in many parts of the world and should be further developed.

## 1. Some world ecosystem questions

Some landscape is perceived by biologists and the public to have value that is real as opposed to financial and is recognized by international conventions, for example, the European Landscape Convention [1]. One valued landscape is upland grazed land, for example in the Pyrenees and other parts of Europe [2]. Such areas are much influenced by farming and very different from the original upland habitat but many would attest to their value. Are they as important as upland woodland? Aspects of the value of land include ecosystem services [3–8] such as water flow regulation, provision of harvested goods, biodiversity preservation and climate stabilization via carbon storage in vegetation and soils. Swetnam *et al.* [8] refer to the value of intact ecosystems, meaning those that are not modified by human activity, but the distinction between modified and unmodified is not always possible or useful. Many of the arguments and quantitative methods developed to calculate value would apply to ecosystems that are partially modified from their original state. Ecosystems subject to some human exploitation can have much biodiversity [9]. The component parts of ecosystems also have value. Populations of charismatic species are of particular interest to the general public, and the lives of the human and non-human individuals present are also valued. Indeed, for many people, the welfare of animals in an area of land is valued more than any other part of the overall system.

Biodiversity is declining in the world, mainly because of farming [10]. Of the total land surface of the world, 33% is used for livestock production [11].

**Table 1.** Reasons for lack of sustainability of a system. (Modified after [32].)

1.	resource depletion	to level that is unacceptable
2.	product accumulation	to level that prevents system function
3.	other effect	to level that people detect and find unacceptable
		to level that affects other systems in an unacceptable way
		to level that affects the system itself, perhaps blocking its function
		to level that is unacceptable
	the consequences of acts or of system functioning (in 1, 2 and 3) could be unacceptable because of immediate or later:	
(a)	harm to the perpetrator	resource loss or poor welfare
(b)	harm to other humans	resource loss
(c)	harm to other humans	poor welfare
(d)	harm to other animals	poor welfare
(e)	harm to environment including that of other animals	

The proportion may well increase in the future so how should livestock areas be managed? Livestock production in Latin America and the Caribbean area has been increasing and today corresponds to 27.1% of the land [12]. Of the 22 million hectares of forest lost between 1960 and 1995, 21 million hectares were then used for cattle production. In tropical regions of the world, annual deforestation rates increased between 2005 and 2010 by 8.5% from an average loss of 10.4 million hectares per year in 2005 [13,14]. Widespread livestock production methods are increasingly viewed as unsustainable, even as the antithesis of conservation [15] and are questioned in relation to animal welfare [16]. One solution, if current widespread animal production systems are used, is to reduce livestock production. Another solution is to use sustainable livestock production methods with much greater on-farm biodiversity than in normal production, no increase in land use and better welfare for the animals [17–19].

The concept of biodiversity includes the extent of variation when the differences considered are genetic [20], biologically functional [21] or based on ecosystem type [22]. Biodiversity may be described numerically [23,24] or by other means [25]. How can adverse effects of livestock production on biodiversity be minimized? Green *et al.* [26] explained that the increase in world food demand, including especially increased demand for animal products, will lead to a reduction in the extent of habitat for wild species of animals and plants and that two solutions for how to reduce this impact have been proposed. One of these is wildlife-friendly farming, whereas another is land sparing and consequent availability of land for nature reserves [27,28]. Green *et al.* produce a model that shows how, to date, farming for species persistence has often depended on demand for agricultural products and how population densities could change with agricultural yield. Land sparing alone leads to islands of ecologically valuable areas in a ‘desert’ of farmland. A combination of land sparing and sustainable farming [29,30] can promote good welfare in animals and much greater *in situ* biodiversity than occurs in the widely used agricultural systems.

## 2. Sustainability

Profitable operation of a system and demand for its products are not sufficient reasons for considering it to be sustainable

and to continue production [31]. Systems were initially called unsustainable when a resource became depleted so much that it became unavailable to the system, or when a product of the system accumulated to a degree that prevented the functioning of the system. Now, the meaning of the term is much wider, for example a system can be unsustainable because of negative impacts on human health, animal welfare or the environment (table 1). Hence, a different definition is required. *A system or procedure is sustainable if it is acceptable now and if its effects will be acceptable in future, in particular in relation to resource availability, consequences of functioning and morality of action* [33,34]. With more criteria for unacceptable harms [32], sustainability is harder to achieve, and unsustainability may be reached long before the production system itself fails. What the public accepts can also change, for example some degree of resource depletion may be tolerated.

Members of the public in all parts of the world, particularly in developed countries, are now insisting on transparency in commercial and governmental activities and on changes in methods of producing of various products. There is a gradual changeover from a ‘push society’, driven in the case of animal production by the producers of the animals, to a ‘pull society’, driven by consumers and facilitated by governments and food retail companies [32,34]. Increasing numbers of consumers now demand ethical production systems and refuse to buy products where production involves, for example, inhumane slaughter methods, rearing calves in small crates, unnecessarily killing dolphins in tuna nets or the payment of very low prices to poor farmers in developing countries. As a consequence, many systems developed with consideration of only short-term market factors, even if widely used at present, are not sustainable. This means that, in some countries, the public have already demanded that such systems do not continue. Throughout the world, the public are likely to make such demands in the relatively near future. The first steps may be the setting up of supply for niche markets, but the rapidity of increase in the consumer pressure is likely to lead to change away from the most unacceptable systems [35]. Changes with small economic cost are likely to occur faster than changes with more substantial cost. One of the first examples of consumers forcing change is the gradual disappearance of animal production procedures with poor welfare for the animals [31–34]. It may be that, in future, consumers will not tolerate very low biodiversity in farmed areas.

### 3. Sustainable silvopastoral production of cattle and other animals

Cattle production for meat, milk or other products may waste valuable resources in that much of the animal feed could be consumed by people or the land used to grow human food plants instead. Additionally, the animals may be kept in such a way that their welfare is poor and, in relation to growth of feed and keeping of animals, there may be adverse impact on the local and world environment. Current cattle production is mainly in large cleared areas in which only herbaceous plants are grown as forage, together with buildings for housing the animals or materials related to production. The effects on the local environment include, initially the removal of trees and shrubs and secondly planting to produce a plant population comprising one, or a very small number of, species. In order to maintain monocultures of pasture plants, herbicides are widely used and biodiversity declines greatly. In addition, there is land used because of construction of roads and buildings, contamination of soil and waterways by agricultural chemicals, carbon cost resulting from CO<sub>2</sub> production from vehicles and from the manufacture of materials used, contamination of water by animal excretions [36], and methane emissions from the animals and their products. In systems, at low or moderate altitude in tropical countries, plant growth rate is relatively fast, and there are often disease and parasite problems. However, modification of cattle production systems to use land resources more effectively, to improve animal welfare and to increase biodiversity concomitant with providing a satisfying and profitable production system [17], is possible in temperate and tropical environments.

A cattle production system is explained here whose characteristics and aims include: using three-level or other multi-level production of edible plants, managing the soil taking account of worms and water retention [11], encouraging predators of harmful animals, minimizing greenhouse gas emissions [37], improving job satisfaction for stock-people, reducing injury and stress in animals and maximizing good welfare, considering how to encourage biodiversity using native shrubs and trees, and using the potential for obtaining wood from trees.

If plant-consuming farm animals, especially ruminants, are fed leaf material, rather than grain, then plant resources otherwise unavailable to humans are used. Although ruminants are of key importance for the sourcing of food for humans in the future, excessive focus on pasture plants for the feeding of farmed ruminants has been a major mistake in almost all parts of the world [38]. Shrubs and trees with edible leaves and shoots, in combination with pasture plants, produce more forage per unit area of land than pasture plants alone. For millennia, trees have often been left in pasture areas, or planted there. Both trees and shrubs can provide shade from hot sun, and shelter from precipitation as well as fulfilling the need of animals to hide from perceived danger [16]. They can also be a substantial source of nutrients for ruminants and other animals [39–43].

Agroforestry has been characterized [44] as being: intentional combination of trees with crops or livestock, intensive in that active management is involved, integrated to enhance the overall productivity of the area and interactive in that the biophysical interactions of component species are manipulated and used. Both selection of plants and management can



**Figure 1.** Cattle browsing *Leucaena* in a silvopastoral system, Caribe, Colombia. (Photograph Walter Galindo, CIPAV).

maximize positive, facilitative interactions among species and minimize negative, competitive interactions [45]. Competition for light can result in a negative impact of trees on pasture plant productivity, in particular if the plants have a C4 photosynthetic pathway with light saturation points at about 85% of full sun. However, pasture plants with a C3 photosynthetic pathway and light saturation photosynthesis at 50% of full sun will not have their growth or yield adversely affected by certain degrees of tree shade [46]. Indeed, shade may improve growth in some pasture plants. Shading increased the protein content, and hence the nutritive value to stock, of the grass *Panicum maximum* in laboratory conditions from 9.6% in Tanzania cultivar plants placed in full sunlight to 12.9% with 54% shading and, in Masai cultivar, 10.5% in sunlight to 15.9% in shading [47]. Nutrient accumulation under woody plants can have long-term beneficial effects for pasture plants [48]. There are complex interactions between foraging ruminants and both plant growth [49] and plant survival [48].

Many of the trees used in agroforestry have leaves and shoots that are toxic or inedible, but, in some cases, the tree species used can provide nutrients for farmed animals, for example, the fruits of shade trees and some 'live fences' [50]. However, it is the planting as forage plants of both shrubs and trees whose leaves and small branches can be consumed by farmed animals that is transforming the prospects for sustainable animal production systems. 'Fodder trees' have been used in several countries, for example, tagasaste *Chamaecytisus palmensis* is widely used by commercial farmers, mainly for cattle feed, in Australia. Individual farmers have pioneered silvopastoral systems, for example in Valle de Cauca, Colombia since 1990. Their feasibility, profitability and consequences for biodiversity have been investigated in detail [51–53].

The key aspect of silvopastoral systems is that the planted food for the animals is not just at the herbaceous level (figure 1). A shrub layer composed of plants that can be eaten by the cattle or other stock is planted and also, in some cases, trees are grown whose leaves can be eaten and any fallen fruits can be consumed. The leaves on the lower branches of trees may be browsed directly or branches can be cut for feeding to stock [54]. Shrubs that are especially suitable for planting as extra food for cattle include *Leucaena leucocephala* and other species of *Leucaena*.

**Table 2.** Changes in nitrogen use and plant production, in an area where cattle were consuming the plants, after a monoculture of a pasture plant *Cynodon plectostachyus* was replaced by the pasture plant plus the leguminous shrub *Leucaena leucocephala*. (Modified after [58].)

variable (per year)	monoculture of <i>Cynodon plectostachyus</i>	silvopastoral system of <i>Leucaena leucocephala</i> (10 000 ha <sup>-1</sup> ) with <i>Cynodon plectostachyus</i>	difference (%)
nitrogen fertilizer ha <sup>-1</sup>	184	0	-100
biomass tonne ha <sup>-1</sup>	23.2	29.9	+29
crude protein tonne ha <sup>-1</sup>	2.5	4.1	+64
metabolizable energy Mcal ha <sup>-1</sup>	56.9	70.2	+23
calcium kg ha <sup>-1</sup>	83.2	142.3	+71
phosphorus kg ha <sup>-1</sup>	74.0	88.8	+20

This leguminous shrub, native to Yucatán in Mexico, has long been used by the Mayans. Recently, it has been used in Northern Australia, Africa, Cuba, other parts of Mexico [55–57], and South America. *Leucaena* is very palatable to cattle, fixates nitrogen, grows very rapidly in tropical conditions and is tolerant of drought [58]. In Colombia, *L. leucocephala* has been planted in intensive silvopastoral systems at a density of 10 000–30 000 shrubs per hectare with pasture plants: *Cynodon plectostachyus*, *Cyperus rotundus*, *Dichanthium aristatum*, *P. maximum* and *Bothriochloa pertusa* [18,59]. Other studies of the combination of trees and pasture plants have been carried out in Veracruz and Jalapa, Mexico [19,60]. A range of tree species, whose leaves and shoots can be eaten by livestock, can be grown in different climatic areas. In the tropics and subtropics of South and Central America, trees that have been used include the leguminous tree *Glracidium sepium* that has high protein, phosphorus, potassium and magnesium in its leaves [61]. Other species with high protein usable by ruminants are *Moringa oleifolia*, in drier areas, *Trichantera gigantea* and *Morus alba* [62].

Because *L. leucocephala* is very palatable to cattle, the animals have to be put in each newly planted area for a short time only, so that they do not damage the plants to the point where they cannot rapidly re-grow. In order to solve this, in silvopastoral systems in Colombia and Mexico a rotational strip system, often separated by electric fences, is used with each group of cattle being moved on every day or every few days. The cattle typically eat the new growth of the *Leucaena* before eating the new grass [18].

If silvopastoral systems are to be advocated, then it is important to consider in what circumstances leaf production that is available for domestic animal consumption can be greater than in pasture-only systems. Table 2 shows, as an example, fodder production in Colombia of a mixed planting of *L. leucocephala* with the grass *C. plectostachyus* in comparison with a monoculture of the pasture plant. Of the material available to cattle, the dry matter production was 27% better, the crude protein production 64% better and the metabolizable energy 23% better in the silvopastoral system [58].

If ruminants are consuming the plants, then their growth and milk production are appropriate measures of the quality of the forage. Milk production in a tropical silvopastoral system, similar to that referred to in the previous paragraph, was 4.13 kg per cow per day when compared with 3.5 kg per day on pasture-only systems. As the numbers of animals per

hectare was much greater, production of good quality milk per hectare was four to five times greater on the silvopastoral system [63]. Milk production from cattle kept on degraded conventional pastures in the humid tropics is very low and, while it can be increased by adding appropriate fertilizer, it can also be greatly improved by silvopastoral system use. Milk production from a silvopastoral system in Colombia, with 4.3 dairy cows per hectare and no artificial fertilizer, was 16 000 l per annum per hectare. The mortality rate was low and the calving interval 12.8 months [18,59]. Milk production can be significantly increased when cattle are able to eat tree leaves in substantial quantities as well as pasture [64]. Milk production on a good quality pasture of the grass *Pennisetum clandestinum* was 12.8 l per cow per day [43]. However, if the cows were also able to eat the leaves of the tree *Alnus acuminata*, then the milk production was 14.4 l per cow per day. In financial terms, the increase in income per cow was from 3152 to 3552 US dollars per cow [63]. If supplements fed to cattle were 75% *Tithonia diversifolia*, a fodder shrub in the family Asteraceae, instead of just conventional concentrates, the milk productivity was increased [18].

#### 4. Soil, nutrients and fertilizer use: silvopastoral and other systems

Much of the structure of the soil is retained in silvopastoral systems with the consequence that earthworms and other soil invertebrates flourish to a greater extent than on land that includes only pasture plants [59]. The presence of deep tree roots, or relatively deep shrub roots, and the maintenance of complex soil structure, has the consequence that water is retained better by the soil in these systems [48]. A further consequence is a reduction in nutrient leaching to ground water. The deep roots of trees and shrubs are capable of retrieving nitrates, and other nutrients that have leached below the rooting zone of herbaceous plants and of eventually recycling these nutrients as litterfall and root turnover in the cropping zone. This role of tree roots has been observed in many cropping systems studied [65]. A silvopastoral system in Florida on flatwood soils (spodosols) was more likely to retain nutrients within plants when compared with plants in an adjacent fertilized pasture with cattle grazing [66]. A comparison in Colombia of soils from conventional pasture for over 30 years and a silvopastoral system for 3, 8 or 12 years found that whereas per cent carbon and

per cent organic matter depended on the amount of clay in the soil, microbial biomass, estimated by total ester-linked fatty acid methylated esters and activities of enzymes such as  $\beta$ -glucosidase, alkaline phosphatase and urease were higher in older silvopastoral areas than in conventional pasture [67]. Conventional pasture favoured Gram-negative bacteria, whereas silvopastoral systems favoured actinomycetes and fungal biomass and there were islands of extra soil fertility under the canopy of trees [68]. Plant growth is thus favoured, as are the production of milk and other animal products.

Silvopastoral systems can result in better conditions for beneficial insects with consequences for soil composition and diversity of plants in the system. Fertilized *C. plectostachyus* pasture and a silvopastoral system with two grasses, *C. plectostachyus* and *P. maximum*, and the shrub *L. leucocephala*, were compared using areas of each system on the same farms in Colombia, and the numbers of dung beetles were higher in the silvopastoral system [66]. There were five species in silvopastoral: three in fertilized, 1.4 times more dung removed per beetle on silvopastoral, 2.7 times as much dung calculated to be removed in total and 1.8 times as many seeds deposited by the beetles. Horn flies, *Hydrotea irritans*, cause irritation to stock and can transmit disease [69]. The numbers trapped on a silvopastoral system were 40% lower than on pasture, probably because of more rapid dung removal and increased numbers of predators of small insects.

The presence of readily degraded manure from the livestock on the pasture and of nitrogen-fixing plants in the silvopastoral system is associated with retention of calcium and phosphorus [58]. One of the advantages of using a nitrogen-fixing shrub species such as *L. leucocephala* is that artificial nitrogenous fertilizers are not required, just supplementary metals in some circumstances [70]. This is a major factor in sustainability as the carbon cost of producing, transporting and applying artificial nitrogen fertilizers is very high.

## 5. Impact of silvopastoral systems on biodiversity and welfare

A remarkable consequence of the use of silvopastoral systems is the increase in biodiversity when compared with pasture-only systems [71,72]. The presence of shrubs and trees very greatly increases cover for wild birds, mammals and reptiles. The greater range of plants increases the number of larger insects, and the more complex soil increases soil insects and other invertebrates. The number of species of birds reported [73] from four areas of silvopastoral systems in Colombia were 108, 135, 137 and 214. The silvopastoral cultivated areas had three times as many bird species as pasture areas without trees in the same region. In another area, there were 24 bird species on pasture without trees, 51 species in woodland and 75 species in silvopastoral systems [74]. Some species in woodland were not present in silvopastoral systems but the impact on biodiversity is clear. There were 30% more ant species on a grass and *Leucaena* system in Colombia, than on grass only, although the major factor affecting ant species numbers was the presence of large trees [75]. Despite these species numbers, there are many plants and animals that are not able to live in silvopastoral farmed land as they require dense forest, extensive marshland or other unmodified habitat in order to survive [76]. For their preservation, separate reserve areas are required.

With increased biodiversity in silvopastoral systems, some of the birds and larger insects whose numbers are increased are predators on ticks, so the numbers of ticks per hectare are reduced, and the prevalence of tick-borne disease reduced. After the implementation of silvopastoral systems and a strategy for the integrated management of ticks, the incidence of anaplasmosis fell from 25% to below 5% in Valle del Cauca, Colombia [77]. In the Cesar region of Colombia, where routine chemical tick control had been required every three weeks, the farms that replaced treeless cattle ranching with silvopastoral systems kept tick numbers low without any chemical tick control [77]. Because tick-borne disease is a major cause of impaired production in the tropical animal production, the impact of the tick predators is of considerable economic importance.

Reduction of ticks, and hence of disease, improves cattle welfare as does reduction of starvation, over-heating and injury [16,78]. In addition to disease reduction, other aspects of poor welfare are also reduced by the presence of shrubs and trees. Starvation is less likely in the silvopastoral systems, which provide a diet with good nutritional composition in dry seasons, than in pasture-only systems. High temperatures can also cause poor welfare, but the shade provided by the shrubs and trees reduces the risk of over-heating. Cattle skin temperatures in a silvopastoral system were 4°C lower than in a pasture-only system [79]. In addition, in some of the systems, the animals have reduced anxiety and fear associated with increased possibility for partial or complete concealment [80]. Even without full concealment, animals in the silvopastoral systems appear more calm and less disturbed by human approach. Such behaviour can be quantitatively evaluated and indicate good welfare. Mancera & Galindo [19], using a range of welfare indicators [81], have shown that the fear response of cattle in areas with more trees is lower than in cows kept in grazing paddocks with fewer trees. They found a reduction in the number of cattle in poor body condition in areas with trees than in areas without trees but with equivalent pasture provision and fewer agonistic interactions than cows with no shade, possibly as a result of the increased stability of the social groups. In a comparison of monoculture and silvopastoral paddocks in Yucatán, Mexico, cattle in the silvopastoral paddocks showed some indication of more cohesive social behaviour and 44% longer resting times. The foraging times were reduced by high temperature and humidity in the monoculture paddocks but not in the silvopastoral paddocks [82].

## 6. Working conditions for stock-people in silvopastoral systems

Sustainability, as defined here, has a human worker component [83]. Workers on silvopastoral farms in Colombia and Mexico, where animal welfare and environmental impact are good, report they like the work and stay in the job longer than people who work on conventional farms [84]. Farmers who adopted silvopastoral systems in the Quindío region of Colombia mentioned different work values as benefits of their new cattle ranching: an increased environmental awareness among workers (29% of employers mentioned it), more initiative and curiosity from employees (21%), a perspective of future job improvement from new knowledge on silvopastoral systems (11%) and reduced social conflict from illegal

**Table 3.** Meat production and carbon emissions in three cattle production systems in Colombia [58] and CIPAV data.

measure	conventional extensive pastures	'improved pastures' without trees	intensive silvopastoral system
animal load (large animals $\text{ha}^{-1}$ )	0.5	1	3
per animal weight gain ( $\text{kg day}^{-1}$ )	0.37	0.5	0.75
per hectare weight gain ( $\text{kg ha}^{-1}$ )	0.185	0.5	2.25
average methane emissions ( $\text{kg ha}^{-1} \text{yr}^{-1}$ )	15.5	38	105
annual meat production – live weight ( $\text{kg ha}^{-1} \text{yr}^{-1}$ )	67.5	182.5	821.3
methane emissions per tonne of meat ( $\text{kg ton}^{-1}$ )	229.5	208.2	127.9
land area required to produce 1 tonne of meat per year (ha)	14.8	5.5	1.2

logging and intrusion (7%) [84]. Several countries have incentive programmes for rural communities, based on payments for ecosystem services.

## 7. Greenhouse gas production and nitrogen usage in silvopastoral systems

The use of shrubs and trees, as well as pasture plants, in animal production systems reduces greenhouse gas production in several ways [58] (table 3). First, carbon loss from growing plants in silvopastoral systems is lower. Second, the loss of carbon from soil is less, because the structure of the soil is maintained better. Third, where trees are browsed, the area is more likely to be used continuously rather than for a short period, so there is less carbon loss when the trees or other plants are removed. Fourth, there is reduced methane production from ruminant animals feeding in the system. Intensive silvopastoral systems produced 12 times more meat than extensive systems, and 4.5 times more meat than 'improved' pastures [85]. Methane emissions increased in a lower proportion: 6.8 and 2.8 times, respectively. Thus, methane emissions per tonne of meat in intensive silvopastoral systems are 1.8 times lower than in extensive cattle ranching. Because three-level production is very efficient in providing food for livestock, less land is required for a given amount of animal production. More production per unit area of land can result in less greenhouse gas emission in the world [18,86].

In a silvopastoral system using hybrid poplar (*Populus* spp.) at a density of 111 trees  $\text{ha}^{-1}$ , the net annual carbon sequestration potential could be as high as  $2.7 \text{ t ha}^{-1} \text{ yr}^{-1}$ , whereas in a monoculture pasture system, the net annual carbon sequestration potential might be less than  $1.0 \text{ t ha}^{-1} \text{ yr}^{-1}$  [87]. Silvopastoral systems with fast-growing tree species therefore have the potential to sequester between two and three times more carbon than monoculture pasture systems. A net annual carbon sequestration rate of  $2.7 \text{ t ha}^{-1} \text{ yr}^{-1}$  is equivalent to an immobilization rate of  $9.9 \text{ t of atmospheric CO}_2 \text{ ha}^{-1} \text{ yr}^{-1}$ . The total carbon sequestered in the permanent woody components of the fast-growing hybrid poplar, together with the carbon contribution to soil from leaf litter and fine root turnover, was approximately  $39 \text{ t C ha}^{-1}$ . Theoretically, this implies that this system has immobilized  $143 \text{ t of CO}_2 \text{ ha}^{-1}$  but 67.5% of the carbon, added via leaf litter and fine roots, was released back

into the atmosphere through microbial decomposition, so the net annual sequestration potential from the trees alone is  $1.7 \text{ t C ha}^{-1} \text{ yr}^{-1}$  or approximately  $6 \text{ t of CO}_2 \text{ ha}^{-1} \text{ yr}^{-1}$  [86]. It has been estimated [88] that carbon-neutrality for the entire Chilean Patagonian cattle industry could be achieved by adopting silvopastoral systems on less than 1% of the total area of the region. However, this calculation might be correct only during rapid tree growth.

Much supplementary nitrogen is typically used in agriculture but most sources are declining and the usage of energy in fertilizer production and in transport of fertilizer to farms is a sustainability factor. *Leucaena leucocephala* and several other shrubs and trees, such as red alder *Alnus rubra* [88], used in silvopastoral systems fix nitrogen to the extent that supplementary nitrogen is usually not required. Future farming systems are more likely to be sustainable if they incorporate nitrogen-fixing plants. For systems involving animal production, there has often been use of rotations with nitrogen-fixing plants grown for only part of the time. There will normally be less nitrogen fixation in such systems than in those in which the nitrogen fixers are present for longer periods, perhaps almost continuously. Indeed, *L. leucocephala* and the mulberry *Morus alba* are sometimes called 'protein banks'. The use of nitrogen-fixing plants native to the area will have the consequence that biodiversity is further increased.

## 8. Temperate and other silvopastoral systems

Forest grazing or browsing in areas managed by humans has long been used in many parts of the world [54]. In oak and pasture systems in the Spanish and French Pyrenees, where many of the trees are *Quercus pyrenaica*, oak leaves may form 25% of the summer diet of goats and 2.5% of that of sheep, whereas both species eat acorns when these become available [89]. Some silvopastoral systems in Portugal, the Mediterranean region and parts of western Asia use planted chestnut trees *Castanea sativa* or *Castanea mollissima*. In many chestnut coppice systems, the major nutrient intake by goats and pigs is understorey plants from April to July, tree leaves from July to October and the fruits of the chestnut from October to December. Chestnut leaves have 12–14% crude protein [89]. Olive (*Olea europaea*) leaves have 12% crude protein and 43% digestible organic matter. Other species of tree used are: *Quercus suber*, *Quercus ilex*, *Alnus nepalensis*, *Sesbania sesbania* and *Pinus radiata*.

## 9. The roles of separate conservation areas and of universal biodiversity increase

Some wildlife can only survive if unmodified forest, marshland, heathland or other natural habitats are available. These habitats have to be of sufficient area for the range used or required by the species. Hence, nature reserves with little or no human modification are required in many parts of the world. However, much of the world, probably an increasing amount [27], will be used for animal production and these areas can be very greatly enriched, in terms of biodiversity, if shrubs and trees as well as pasture plants are present. If they form wildlife corridors, for example along water-courses, then their impact on world biodiversity is likely to be increased [66]. Although people are unlikely to pay to see pasture without trees, the greatly increased numbers of birds and other wildlife in silvopastoral systems with trees may offer economic opportunity for ecotourism. A combination of nature reserves and large areas of species-rich systems, in which the welfare of the animals produced is good, is likely to be demanded by an increasing proportion of the public in all parts of the world.

## 10. System uptake by farmers

Are silvopastoral systems likely to be taken up by farmers? Agroforestry methods and new forage plants have often not been readily used by farmers. Some systems have not spread, because a financial return takes 3–6 years [90]. Lack of security of land tenure may also deter farmers from investing in future yields. However, the planting of *Leucaena* as part of a silvopastoral system can lead to substantial forage availability within nine months in tropical conditions. A further cost of some innovative changes in forage plants is that of maintaining the plant

system. The nitrogen-fixing plant has to grow well in competition with pasture plants and *Leucaena* certainly does so. Palatable shrubs have to be protected from destruction by grazing or browsing animals, for example by limiting time in the forage area. There is some cost associated with moving animals, and electric fences but extra plant production compensates for this. There seems to be increasing usage of silvopastoral systems in several tropical and temperate countries.

## 11. Conclusion

Animal protein from herbivorous mammals is important for providing human food. When ruminants are farmed, and they are fed materials that cannot be digested by humans, such as leaves and other cellulose-containing tissue, there is a positive net effect on human food provision. However, can ruminant production systems be sustainable? A system or procedure is sustainable if it is acceptable now and if its effects will be acceptable in future, in particular in relation to resource availability, consequences of functioning and morality of action. The advantages of silvopastoral systems for increasing biodiversity, improving animal welfare, providing good working conditions and allowing a profitable farming business are such that these systems are sustainable where many other large herbivore production systems are not. With good management, silvopastoral systems can replace existing systems in many parts of the world, reducing agricultural expansion into conservation areas. There should be further work developing them [91].

**Acknowledgements.** We thank Zoraida Calle, David Williams, other colleagues in our three institutes and the referees of this paper for helpful advice.

**Funding statement.** We thank PAPIIT-UNAM for financial support.

## References

- Council of Europe. 2000 *European landscape convention*, pp. 176. Florence, Italy: European Treaty Series.
- Gibon A. 2005 Managing grassland for production, the environment and the landscape. Challenges at the farm and the landscape level. *Livest. Prod. Sci.* **96**, 11–31. (doi:10.1016/j.livprodsci.2005.05.009)
- Costanza R *et al.* 1997 The value of the world's ecosystem services and natural capital. *Nature* **387**, 253–260. (doi:10.1038/387253a0)
- de Groot R, Wilson MA, Boumans RM. 2002 A typology for the classification, description and valuation of ecosystem functions, goods and services. *Ecol. Econ.* **41**, 393–408. (doi:10.1016/S0921-8009(02)00089-7)
- Boyd J, Banzhaf S. 2007 What are ecosystem services? The need for standardized environmental accounting units. *Ecol. Econ.* **63**, 616–626. (doi:10.1016/j.ecolecon.2007.01.002)
- Ruhl JB, Kraft SE, Lant CL. 2007 *The law and policy of ecosystem services*. Washington, DC: Island Press.
- Berber P, Mooney HA, Dijkman J, Tarawali S, de Haan C (eds) 2010 *Livestock in a changing landscape: experiences and regional perspectives*, vol. 2. Washington, DC: Island Press.
- Swetnam RD *et al.* 2011 Mapping socio-economic scenarios of land cover change: a GIS method to enable ecosystem service modelling. *J. Environ. Manage.* **92**, 563–574. (doi:10.1016/j.jenvman.2010.09.007)
- Kull CA, Carrière SM, Moreau S, Ramiarantsoa HR, Blanc-Pamard C, Tassin J. 2013 Melting pots of biodiversity: tropical smallholder farm landscapes as guarantors of sustainability. *Environ. Sci. Policy Sustain. Dev.* **55**, 6–16. (doi:10.1080/00139157.2013.765307)
- CBD, Secretariat of the Convention on Biological Diversity. 2006 *Global biodiversity outlook*, 2. Montreal, Canada: World Trade Centre.
- Giraldo C, Escobar F, Chará JA, Calle Z. 2011 The adoption of silvopastoral systems promotes the recovery of ecological processes regulated by dung beetles in the Colombian Andes. *Insect Conserv. Divers.* **4**, 115–122. (doi:10.1111/j.1752-4598.2010.00112.x)
- FAO. 2009 *The state of world agriculture and food 2009. Cattle production examined*. Rome, Italy: FAO.
- FAO. 2006 *Global forest resources assessment 2005*. FAO Forestry Paper 147, Rome, Italy: FAO.
- Jarvis A, Touval JL, Castro M, Sotomayor L, Hyman GG. 2010 Assessment of threats to ecosystems in South America. *J. Nat. Conserv.* **18**, 180–188. (doi:10.1016/j.jnc.2009.08.003)
- Perfecto I, Vandermeer J. 2010 Biodiversity conservation in tropical agroecosystems: a new conservation paradigm. *Proc. Natl Acad. Sci. USA* **107**, 5786–5791. (doi:10.1073/pnas.0905455107)
- Broom DM, Fraser AF. 2007 *Domestic animal behaviour and welfare*, 4th edn. Wallingford, UK: CABI.
- Murgueitio E, Cuartas CA, Naranjo JF. 2008 *Ganadería del Futuro*. Cali, Colombia: Fundación CIPAV.
- Murgueitio E, Calle Z, Uribe F, Calle A, Solorio B. 2011 Native trees and shrubs for the productive rehabilitation of tropical cattle ranching lands. *Forest Ecol. Manage.* **261**, 1654–1663. (doi:10.1016/j.foreco.2010.09.027)
- Mancera AK, Galindo F. 2011 *Evaluation of some sustainability indicators in extensive bovine*

- stockbreeding systems in the state of Veracruz.* VI Reunión Nacional de Innovación Forestal 31. México: León Guanajuato.
20. Vane-Wright R. 1991 What to protect? – systematics and the agony of choice. *Biol. Conserv.* **55**, 235–254. (doi:10.1016/0006-3207(91)90030-D)
  21. Petachy O, Gaston K. 2002 Functional diversity (FD), species richness and community composition. *Ecol. Lett.* **5**, 402–411. (doi:10.1046/j.1461-0248.2002.00339.x)
  22. Weikard H. 2002 Diversity functions and the value of biodiversity. *Land Econ.* **78**, 20–27. (doi:10.2307/3146920)
  23. Jost L. 2006 Entropy and diversity. *Oikos* **113**, 363–374. (doi:10.1111/j.2006.0030-1299.14714.x)
  24. Jost L. 2008  $G_{ST}$  and its relatives do not measure differentiation. *Mol. Ecol.* **17**, 4015–4026. (doi:10.1111/j.1365-294X.2008.03887.x)
  25. Hoffmann S, Hoffmann A. 2008 Is there a ‘true’ diversity? *Ecol. Econ.* **65**, 213–213. (doi:10.1016/j.ecolecon.2008.01.009)
  26. Green RE, Cornell SJ, Scharlemann JPW, Balmford A. 2005 Farming and the fate of wild nature. *Science* **307**, 550–555. (doi:10.1126/science.1106049)
  27. Balmford A, Green R, Phalan B. 2012 What conservationists need to know about farming. *Proc. R. Soc. B* **279**, 2714–2724. (doi:10.1098/rspb.2012.0515)
  28. Tilman D, Cassman KG, Matson PA, Naylor R, Polasky S. 2002 Agricultural sustainability and intensive production practices. *Nature* **418**, 671–677. (doi:10.1038/nature01014)
  29. Daily GC, Ceballos G, Pacheco J, Suzan G, Sanchez-Azofeifa A. 2003 Countryside biogeography of neotropical mammals: conservation opportunities in agricultural landscapes of Costa Rica. *Conserv. Biol.* **17**, 1814–1826. (doi:10.1111/j.1523-1739.2003.00298.x)
  30. Scherr SJ, McNeely JA. 2007 *Farming with nature*. Washington, DC: Island Press.
  31. Aland A, Madec F (eds) 2009 *Sustainable animal production*, pp. 496. Wageningen, The Netherlands: Wageningen Academic Publishers.
  32. Broom DM. 2012 Defining agricultural animal welfare: from a sustainability and product quality viewpoint. In *Animal welfare in animal agriculture* (eds WG Pond, FW Bazer, BE Rollin), pp. 84–91. Boca Raton, FL: CRC Press.
  33. Broom DM. 2001 The use of the concept animal welfare in European conventions, regulations and directives. In *Food chain*, pp. 148–151. Uppsala, Sweden: SLU Services, Uppsala.
  34. Broom DM. 2010 Animal welfare: an aspect of care, sustainability, and food quality required by the public. *J. Vet. Med. Educ.* **37**, 83–88. (doi:10.3138/jvme.37.1.83)
  35. Pollan M. 2006 *The omnivore’s dilemma: a natural history of four meals*. New York, NY: Penguin Books.
  36. Chará JD, Pedraza GX, Giraldo LP. 2008 Protección de ambientes acuáticas en zonas ganaderas. In *Ganadería del Futuro* (eds E Murgeutio, CA Cuartas, JF Naranjo), pp. 111–129. Cali, Colombia: Fundación CIPAV.
  37. Naranjo JF, Cuartas CA, Murgeutio E, Chará J, Barahona R. 2012 Balance de gases de efecto invernadero en sistemas silvopastoriles intensivos con *Leucaena leucocephala* en Colombia. *Livest. Res. Rur. Dev.* **24**, 150.
  38. Preston TR, Leng RA. 1987 *Matching ruminant production systems with available resources in the tropics and sub-tropics*. Armidale, Australia: Penambul Books.
  39. Harvey CA, Haber WA. 1998 Remnant trees and the conservation of biodiversity in Costa Rican pastures. *Agroforest. Syst.* **44**, 37–68. (doi:10.1023/A:1006122211692)
  40. Souza de Abreu MH, Ibrahim M, Harvey C, Jiménez F. 2000 Caracterización del componente arbóreo en los sistemas en la Fortuna de San Carlos, Costa Rica. *Agriforestaria de las Américas* **7**, 53–56. (Turrialba, Costa Rica, CATIE).
  41. Esquivel H, Ibrahim M, Harvey C, Villanueva G. 2004 Árboles dispersos en potreros da fincas ganaderas en un ecosistema seco da Costa Rica. In *Agriforestaria de las Américas*, vol. 11. Turrialba, Costa Rica: CATIE.
  42. Fornos S, Espinel RG, Murgeutio E, Mejia C, Soto R, García B. 2008 Establecimiento y desarrollo de sistemas silvopastoriles en Nicaragua. In *Ganadería del Futuro* (eds E Murgeutio, CA Cuartas, JF Naranjo), pp. 301–323. Cali, Colombia: Fundación CIPAV.
  43. Chamorro D, Rey AM. 2008 El componente arbóreo como dinamizador del sistema de producción de leche en el trópico alto Colombiano, experiencias de Corpoica-Tibaitatá. In *Ganadería del Futuro* (eds E Murgeutio, CA Cuartas, JF Naranjo), pp. 349–397. Cali, Colombia: Fundación CIPAV.
  44. Gold MA, Garrett HE. 2009 Agroforestry nomenclature, concepts, and practices. In *North American agroforestry: an integrated science and practice* (ed. HE Garrett), 2nd edn. Madison, WI: Agronomy Society of America.
  45. Jose S, Gillespie AR, Pallardy SG. 2004 Interspecific interactions in temperate agroforestry. *Agroforest. Syst.* **61**, 237–255. (doi:10.1023/B:AGFO.0000029002.85273.9b)
  46. Sharow SH. 1999 Silvopastoralism: competition and facilitation between trees, livestock, and improved grass-clover pastures on temperate rainfed lands. In *Agroforestry in sustainable agricultural systems* (eds LE Buck, I Lassoie, ECM Fernandez), pp. 111–130. Boca Raton, FL: CRC Press.
  47. Silveira SR et al. 2012 Valor nutritivo de *Panicum maximum*, cultivares Tanzania y Massai, de acuerdo a niveles de sombra y de fertilización nitrogenada. In *Proc. 2nd Congreso Nacional de Sistemas Silvopastoriles* (eds J Arroquy, R Ledesma, B Roldán, A Gómez), pp. 289–293. Santiago del Estero, Argentina: INTA, Min Agric.
  48. Scholes RJ, Archer SR. 1997 Tree-grass interactions in savannas. *Annu. Rev. Ecol. Syst.* **28**, 517–544. (doi:10.1146/annurev.ecolsys.28.1.517)
  49. Broom DM, Arnold GW. 1986 Selection by grazing sheep of pasture plants at low herbage availability and responses of the plants to grazing.
  50. Aust. J. Agric. Res. **37**, 527–538. (doi:10.1071/AR9860527)
  51. Harvey CA et al. 2005 Contribution of live fences to the ecological integrity of agricultural landscapes in Central America. *Agric. Ecosyst. Environ.* **111**, 200–230. (doi:10.1016/j.agee.2005.06.011)
  52. Murgeutio E, Calle Z. 1999 Diversidad biológica en sistemas de ganadería bovina en Colombia. In *Agroforestería para la producción animal en Latinoamerica. Estudio FAO sobre producción y sanidad animal*, vol. 143 (eds M Sánchez, M Rosales). Rome, Italy: FAO.
  53. Murgeutio E. 2000 Sistemas agrororestales de la producción ganadera en Colombia. In *Intensificación de la Ganadería en Centroamérica – Beneficios, Económicos y Ambientales* (eds C Pomareda, H Steinfield), pp. 219–242. San José, Costa Rica: CATIE, FAO and SIDE.
  54. Murgeutio E. 2004 Silvopastoral systems in the neotropics. In *Proc. Int. Conf. on Silvopastoralism and Sustainable Management*, Lugo, Spain (eds MR Mosquera-Losada, A Rigueiro-Rodriguez, J McAdam), pp. 24–29. Lugo, Wallingford, UK: CABI.
  55. Etienne M. 1996 *Temperate and Mediterranean silvopastoral systems of Western Europe*. Paris, France: INRA.
  56. National Academy of Sciences. 1977 *Leucaena promising forage and tree crop for the tropics*. Washington, DC: National Academy of Sciences.
  57. Dalzell DA, Shelton HM, Mullen BF, Larsen PH, McLaughlin KG. 2006 *Leucaena: a guide to establishment and management*. Sydney, Australia: Meat and Livestock Australia Ltd.
  58. Flores M, Solorio B. 2012 *Ganadería sustentable en México, primera etapa del proyecto estratégico de calidad nacional*. SAGARPA, Fundación Produce Michoacán, 47 p. Morelia, Michoacán, México: COFUPRO, UADY.
  59. Molina CCH, Molina DCH, Molina EJ, Molina JP. 2008 Carne, leche y medio ambiente en el sistema silvopastoril intensivo con *Leucaena leucocephala* (Lam.) de Wit Mimosaceae. In *Ganadería del Futuro* (eds E Murgeutio, CA Cuartas, JF Naranjo), pp. 41–65. Cali, Colombia: Fundación CIPAV.
  60. Solorio-Sánchez FJ, Solorio-Sánchez B, Casanova-Lugo F, Ramírez-Avilés L, Ayala-Burgos A, Ku-Ver J, Aguilar-Pérez C. 2012 Situación actual global de la investigación y desarrollo tecnológico en el establecimiento, manejo e aprovechamiento de los sistemas silvopastoriles intensivos. In *Memorias del IV Congreso Internacional sobre Sistemas Silvopastoriles*. Morelia, Michoacán, Mexico.
  61. Stewart JL, Allison GE, Simons AJ. 1996 *Gliricidia sepium: genetic resources for farmers*. Oxford, UK: Oxford Forestry Institute.
  62. Leterme P, Bulgen A, Murgeutio E, Cuartas C. 2007 *Fodder banks for sustainable pig production systems*. Cali, Colombia: CIPAV.

63. Murgueitio E, Chará J, Uribe F, Gacharná N, Tafur 2011 Pesquisa, desenvolvimento e inovação para la sustentabilidade da bovinocultura leiteira. In *Juiz da Fora (MG)* (eds LG Ribeiro Pereira), pp. 93–108. Brasil: EMBRAPA Gado de leite.
64. Flores M, Solorio SB. 2011 Proyecto estratégico de prioridad nacional para el establecimiento de sistemas silvopastoriles intensivos en temporados de lluvias y sequías en la región del Caribe de Colombia. In *Memorias del Tercer Congreso sobre Sistemas Silvopastoriles para la Ganadería Sostenible del siglo XXI*, Morelia, Michoacán, México, pp. 32–45. Cali, Colombia: CIPAV.
65. Allen SC, Jose S, Nair PKR, Brecke BJ, Nkedi-Kizza P, Ramsey CL. 2004 Safety-net role of tree roots: evidence from a pecan (*Carya illinoensis* K. Koch)–cotton (*Gossypium hirsutum* L.) alley cropping system in the southern United States. *Forest Ecol. Manage.* **192**, 395–407. (doi:10.1016/j.foreco.2004.02.009)
66. Nair VD, Graetz DA. 2004 Agroforestry as an approach to minimizing nutrient loss from heavily fertilized soils: the Florida experience. *Agroforest. Syst.* **61**, 269–279. (doi:10.1023/B:AGFO.0000029004.03475.1d)
67. Vallejo VD, Roldan F, Dick RP. 2010 Soil enzymatic activities and microbial biomass in an integrated agroforestry biosequence compared to monoculture and a native forest of Colombia. *Biol. Fertil. Soils* **46**, 577–587. (doi:10.1007/s00374-010-0466-8)
68. Vallejo VD, Arbeli Z, Terán W, Lorenz N, Dick RP, Roldan F. 2012 Effect of land management and *Prosopis juliflora* (Sw.) DC trees on soil microbial community and enzymatic communities in intensive silvopastoral systems of Colombia. *Agric. Ecosyst. Environ.* **150**, 139–148. (doi:10.1016/j.agee.2012.01.022)
69. Hillerton JE, Bramley AJ, Broom DM. 1983 *Hydrotaea irritans* and summer mastitis in calves. *Vet. Rec.* **113**, 88–89. (doi:10.1136/vr.113.4.88)
70. Ceballos A, Rubio V, Estrada J. 2008 Desequilibrios minerales de bovinos en sistemas silvopastoriles. In *Ganadería del Futuro* (eds E Murgueitio, CA Cuartas, JF Naranjo), pp. 431–439. Cali, Colombia: Fundación CIPAV.
71. Burgess PJ. 1999 Effects of agroforestry on farm biodiversity in the UK. *Scottish Forestry* **53**, 24–27.
72. McAdam JH, McEvoy PM. 2009 The potential for silvopastoralism to enhance biodiversity on grassland farms in Ireland. In *Agroforestry in Europe: current status and future prospects* (eds A Rigueiro-Rodriguez, J McAdam, MR Mosquera-Losada), pp. 343–356. Berlin, Germany: Springer.
73. Fajardo ND, Johnston R, Neira LA. 2008 Sistemas ganaderos amigos de los aves. In *Ganadería del Futuro* (eds E Murgueitio, CA Cuartas, JF Naranjo), pp. 171–203. Cali, Colombia: Fundación CIPAV.
74. Múnera E, Bock BC, Bolívar DM, Botero JA. 2008 Composición y estructura de la avifauna en diferentes hábitats en el departamento de Córdoba, Colombia. In *Ganadería del Futuro* (eds E Murgueitio, CA Cuartas, JF Naranjo), pp. 205–225. Cali, Colombia: Fundación CIPAV.
75. Rivera LB, Botero M, Escobar S, Armbrrecht I. 2008 Diversidad de hormigas en sistemas ganaderos. In *Ganadería del Futuro* (eds E Murgueitio, CA Cuartas, JF Naranjo), pp. 228–243. Cali, Colombia: Fundación CIPAV.
76. Phalan B, Balmford A, Green RE, Scharlemann JPW. 2011 Minimising the harm to biodiversity of producing more food globally. *Food Policy* **26**, 562–571.
77. Murgueitio E, Giraldo C. 2009 Sistemas silvopastoriles y control de parásitos. *Revista Carta Fedegán* **115**, 60–63.
78. Petherick JC. 2005 Animal welfare issues associated with extensive livestock production: the northern Australian beef cattle industry. *Appl. Anim. Behav. Sci.* **92**, 211–234. (doi:10.1016/j.applanim.2005.05.009)
79. Galindo F, Olea R, Suzán G. 2013 Animal welfare and sustainability. International Workshop on Farm Animal Welfare. São Paulo SP Brazil. <http://www.workshopdebemestanimal.com.br/indexen.html>.
80. Ocampo A, Cardozo A, Tarazona A, Ceballos M, Murgueitio E. 2011 La investigación participativa en bienestar y comportamiento animal en el trópico de América: oportunidades para nuevo conocimiento aplicada. *Revista Colombiana Ciencias Pecuarias* **24**, 332–346.
81. Blokhuis HJ, Jones RB, Geers R, Miele M, Veissier I. 2003 Measuring and monitoring animal welfare: transparency in the food product quality chain. *Anim. Welfare* **12**, 445–455.
82. Améndola L, Solorio FJ, González-Rebeles C, Galindo F. 2013 Behavioural indicators of cattle welfare in silvopastoral systems in the tropics of México. In *Proc. 47th Congress Int. Soc Appl Ethol*, Florianópolis, 150.
83. Viljoen AJ, Wiskerke SC (eds). 2012 *Sustainable food planning: evolving theory and practice*. Wageningen, The Netherlands: Wageningen Academic Publishers.
84. Calle A, Montagnini F, Zuluaga AF. 2009 Farmers' perceptions of silvopastoral system production in Quindío, Colombia. *Bois et forêts des tropiques* **300**, 79–94.
85. Murgueitio E, Chará J, Barahona R, Cuartas CA, Naranjo JF. 2012 Los sistemas silvopastoriles intensivos, herramienta da mitigación y adaptación del cambio climático. *Proc IV Congreso Internacional Sistemas Silvopastoriles Intensivos*, 1–8. Morelia y Valle de Apatzingán/Tepalcatepec. Fundación Produce Michoacán, Cofupro, Sagarpa, CIPAV, UADY. Morelia (Michoacán), México.
86. Burney JA, Davis SJ, Lobell DB. 2010 Greenhouse gas mitigation by agricultural intensification. *Proc. Natl Acad. Sci. USA* **107**, 12 052–12 057. (doi:10.1073/pnas.0914216107)
87. Gordon AM, Thevathasan NV, Dube F. 2012 Carbon sequestration: a farm-level benefit of adopting silvopastoralism in temperate North America. In *Proc. 2nd Congreso Nacional de Sistemas Silvopastoriles* (eds J Arroquy, R Ledesma, B Roldán, A Gómez), pp. 429–440. INTA, Min Agric. Santiago del Estero: Argentina Santiago del Estero.
88. Dube F, Zagal EN, Stolpe M, Espinosa. 2009 The influence of land-use change on the organic carbon distribution and microbial respiration in a volcanic soil of the Chilean Patagonia. *Forest Ecol. Manage.* **257**, 1695–1704. (doi:10.1016/j.foreco.2009.01.044)
89. Castro M, Castro JF, Gómez-Sal A. 2004 *Quercus pyrenaica* Willd. Woodlots and small ruminant production in North east Portugal. In *Sustainability of agro-silvopastoral systems, dehesas, montados* (eds S Schnabel, A Ferreira), chapter 5. Reiskirchen, Germany: Catena.
90. Mercer DE. 2004 Adoptions of agroforestry innovations in the tropics: a review. *Agroforest. Syst.* **61–62**, 311–328. (doi:10.1023/B:AGFO.0000029007.85754.70)
91. Teklehaimanot Z, Mmolotsi RM. 2007 Contribution of red alder to soil nitrogen input in silvopastoral systems. *Biol. Fertil. Soils* **43**, 843–848. (doi:10.1007/s00374-006-0163-9)